

PERIODIC SOLUTIONS OF SINGULAR INTEGRAL EQUATIONS

T. A. Burton

Northwest Research Institute
732 Caroline Street
Port Angeles, WA 98362
taburton@olympen.com

Bo Zhang

Department of Mathematics and Computer Science
Fayetteville State University
Fayetteville, NC 28301
bzhang@uncfsu.edu

ABSTRACT. We consider a scalar integral equation $x(t) = a(t) - \int_{-\infty}^t C(t, s)g(s, x(s))ds$ in which $C(t, s)$ has a singularity at $t = s$. There are periodic assumptions on a , C , and g . First we prove a fixed point theorem of the Krasnoselskii-Schaefer type. We then construct a Liapunov functional which allows us to satisfy the conditions of the fixed point theorem and to prove that there is a periodic solution.

1. INTRODUCTION

We consider a scalar integral equation

$$(1) \quad x(t) = a(t) - \int_{-\infty}^t C(t, s)g(s, x(s))ds$$

for which there is a $T > 0$ so that

$$(2) \quad a(t+T) = a(t), g(t+T, x) = g(t, x), C(t+T, s+T) = C(t, s)$$

for all $t \in \mathfrak{R}$ and $s < t$ with a and g continuous. We denote by $(\mathcal{P}_T, \|\cdot\|)$ the Banach space of continuous T -periodic functions.

If g is Lipschitz and if C is small enough then a contraction mapping will yield a periodic solution. If C is convex then Liapunov arguments will produce *a priori* bounds. Under compactness conditions, Schaefer's fixed point theorem will yield a periodic solution. A collection of such results are found in Burton [6]. A recent n -dimensional result is given in [16].

In this paper we ask that g satisfies

$$(3) \quad |g(t, x) - g(t, y)| \leq K|x - y|$$

2000 Mathematics Subject classifications: 45D05, 45D20, 45M15

Key words: Integral Equations, Fixed Point Theorems, Boundedness, Periodic Solutions, Liapunov Functions

for all $x, y \in \mathfrak{R}$ and some $K > 0$, while C satisfies a truncated convexity condition, but has a significant singularity at $t = s$. We derive a set of conditions measuring the magnitude of the singularity that will still permit proof of the existence of a periodic solution using a combination Krasnoselskii-Schaefer fixed point theorem which we will prove in Section 2.

2. A FIXED POINT THEOREM

In this section, we will prove a fixed point theorem of Krasnoselskii-Schaefer type in which the mapping function has the form $Px = Bx + Ax$ with A being compact and $(I - B)^{-1}$ continuous on an appropriate subset M of a Banach space S . The theorem resembles that of Burton-Kirk [5] without having a λ term in B . See [7], [9], [10], [12], [13], [14] for work on Krasnoselskii and Schaefer theorems and their extended forms.

Since P is the sum of two operators, it is in general a non-self map; that is, P may not necessarily map a closed convex subset M of S into itself. To prove the existence of a fixed point of P , we apply topological degree theory or transversality method by constructing a homotopy U_λ on M with $U_1 = P$. It is assumed that $U_\lambda(\phi) = U(\lambda, \phi)$ is a continuous mapping of $[0, 1] \times M$ into a compact subset of S . In many applications, U_0 is a constant map sending M to a point $p \in M/\partial M$. In this case, U_0 is an "essential" map. If $U_\lambda(\phi)$ is fixed point free on ∂M for all $\lambda \in (0, 1)$, then $U_1(\phi)$ is essential having a fixed point property in M (Granas and Dugundji [8, p.120-123]). This fact is often written in the form of Leray-Schauder Principle or its nonlinear alternatives which states that either

- (A₁) U_1 has a fixed point in M or
- (A₂) there exists $x \in \partial M$ and $\lambda \in (0, 1)$ with $x = U_\lambda(x)$

(see [1, p.48], [8, p.123], [14, p.28], [15]).

Theorem 2.1. *Let $(S, \|\cdot\|)$ be a Banach space, $A, B : S \rightarrow S$ such that A is continuous with A mapping bounded sets into compact sets, $(I - B)^{-1}$ exists and is continuous on $(I - B)S$ with $\lambda A(M) \subset (I - B)S$ for each closed convex subset $M \subset S$ and $\lambda \in [0, 1]$. Then either*

- (i) $x = Bx + \lambda Ax$ has a solution in S for $\lambda = 1$, or
- (ii) the set of all such solutions, $0 < \lambda < 1$, is unbounded.

Proof. Since $\lambda A(M) \subset (I - B)S$, we have $0 \in (I - B)S$. If $x^* = (I - B)^{-1}(0)$, then x^* is the unique fixed point of B . For each positive integer n , define a closed and bounded set

$$M_n = \{x \in S : \|x\| \leq n\}.$$

We choose n sufficiently large so that $x^* \in M_n/\partial M_n$. Now $(I - B)^{-1}$ exists and is continuous on $(I - B)S$. Since A is continuous with A mapping M_n into a compact set, so is $(I - B)^{-1}(\lambda A)$ for each $\lambda \in [0, 1]$. Define $U : [0, 1] \times M_n \rightarrow S$ by

$$U(\lambda, \phi) = (I - B)^{-1}(\lambda A\phi).$$

Then $U_\lambda(\phi) = U(\lambda, \phi)$ is a continuous mapping of $[0, 1] \times M_n$ into a compact subset of S . Indeed, set $\Gamma = \{\lambda A\phi : \lambda \in [0, 1], \phi \in M_n\}$ and let $\{(\lambda_k, \phi_k)\}$ be a sequence in $[0, 1] \times M_n$. We may assume that $\lambda_k \rightarrow \lambda_0 \in [0, 1]$ as $k \rightarrow \infty$. Since AM_n is contained in a compact subset of S , there exists a convergent subsequence $\{A\phi_{k_j}\}$

of $\{A\phi_k\}$. Now $\{\lambda_{k_j}A\phi_{k_j}\}$ converges in S . This implies that Γ is pre-compact, and so is $(I - B)^{-1}\Gamma$. Observe that for all $\phi \in M_n$,

$$U_0(\phi) = (I - B)^{-1}(0) = x^*$$

is a constant map. Moreover, $x^* \in M_n/\partial M_n$. By the statement of nonlinear alternatives (A₁) and (A₂) above, either U_1 has a fixed point in M_n or there exists $x_n \in \partial M_n$ such that $x_n = U_\lambda(x_n)$ for some $\lambda \in (0, 1)$. This implies that either $x = Bx + Ax$ has a solution in M_n or there exists $x_n \in \partial M_n$ with $x_n = Bx_n + \lambda Ax_n$ for some $\lambda \in (0, 1)$. In the later case, we have $\|x_n\| = n$. Thus, if (i) does not hold, then $\|x_n\| \rightarrow \infty$ as $n \rightarrow \infty$ and (ii) must hold. This completes the proof. \square

Remark 2.2. It is clear that if B is a contraction mapping with contraction constant $0 < \alpha < 1$, then $(I - B)^{-1}$ exists and is continuous on S . Many generalized or nonlinear contractions satisfy this condition (see [2], [3], [7], [10], [11], [12]).

3. TECHNICAL CONDITIONS

We now introduce the conditions which will produce the *a priori* bound needed in the fixed point theorem, as well as the required compactness. The kernel, $C(t, s)$, can have a singularity at $t = s$, but we ask that there exist a fixed $\epsilon > 0$ so that

$$(4) \quad C(t, s) \geq 0, C_s(t, s) \geq 0, C_t(t, s) \leq 0, C_{st}(t, s) \leq 0$$

provided that

$$(5) \quad -\infty < s \leq t - \epsilon, t < \infty.$$

Moreover, if $x \in \mathcal{P}_T$, then

$$(6) \quad \int_{-\infty}^{t-\epsilon} C(t, s)g(s, x(s))ds \text{ and } \int_{t-\epsilon}^t C(t, s)g(s, x(s))ds \text{ are continuous.}$$

The ϵ will play a central role. First, assume that there is a $\eta < 1$ with

$$(7) \quad K \int_{t-\epsilon}^t |C(t, s)|ds \leq \eta, t \in \mathfrak{R}.$$

Next, there are positive constants α and β with $2\alpha + \beta < 2$ so that both

$$(8) \quad \int_s^{s+\epsilon} [\epsilon C_s(u, u - \epsilon) + C(u, u - \epsilon) + |C(u, s)|]du < \alpha, s \in \mathfrak{R}$$

and

$$(9) \quad C(t, t - \epsilon)\epsilon + \int_{t-\epsilon}^t |C(t, s)|ds < \beta, t \in \mathfrak{R}.$$

The work here is motivated by and is an extension of [4]. Relations (7)–(9) specify the strength of the singularity. For a “mild” singularity such as $C(t, s) = [t - s]^{-p}$, $0 < p < 1$, then (4), (5), (7)–(9) are satisfied for any $K > 0$ when it is allowed that ϵ can be taken sufficiently small. But (6) would fail. The following function satisfies (4)–(9) with $0 < \epsilon \leq 1$ and an appropriate constant $k > 0$

$$C(t, s) = \frac{k}{(t - s)(1 + |\ln(t - s) - \ln \epsilon|)^2}.$$

We now define for $0 \leq \lambda \leq 1$ a companion equation to (1)

$$(1\lambda) \quad x(t) = \lambda \left[a(t) - \int_{-\infty}^{t-\epsilon} C(t, s)g(s, x(s))ds \right] - \int_{t-\epsilon}^t C(t, s)g(s, x(s))ds.$$

The mappings $A, B : \mathcal{P}_T \rightarrow \mathcal{P}_T$ mentioned in the theorem are defined by $\phi \in \mathcal{P}_T$ implies that

$$(10) \quad (A\phi)(t) := a(t) - \int_{-\infty}^{t-\epsilon} C(t, s)g(s, \phi(s))ds$$

and

$$(11) \quad (B\phi)(t) := - \int_{t-\epsilon}^t C(t, s)g(s, \phi(s))ds.$$

By (6), if $\phi \in \mathcal{P}_T$ then ϕ is continuous so these integrals are continuous functions. To see that $A\phi, B\phi \in \mathcal{P}_T$ we note that

$$\begin{aligned} (A\phi)(t+T) &= a(t+T) - \int_{-\infty}^{t+T-\epsilon} C(t+T, s)g(s, \phi(s))ds \\ &= a(t) - \int_{-\infty}^{t-\epsilon} C(t+T, s+T)g(s+T, \phi(s+T))ds = (A\phi)(t) \end{aligned}$$

while

$$\begin{aligned} (B\phi)(t+T) &= - \int_{t+T-\epsilon}^{t+T} C(t+T, s)g(s, \phi(s))ds \\ &= - \int_{t-\epsilon}^t C(t+T, s+T)g(s+T, \phi(s+T))ds = (B\phi)(t). \end{aligned}$$

Moreover, by (3) and (7), B is a contraction.

4. A LIAPUNOV FUNCTIONAL

We begin with the assumption that there is an $L > 0$ with

$$(12) \quad xg(t, x) \geq 0 \text{ for } |x| \geq L$$

and that

$$(13) \quad \lim_{s \rightarrow -\infty} (t-s)C(t, s) = 0 \text{ for fixed } t.$$

Then define a Liapunov functional by

$$(14) \quad V(t, \epsilon) = \lambda \int_{-\infty}^{t-\epsilon} C_s(t, s) \left(\int_s^t g(v, x(v))dv \right)^2 ds.$$

Lemma 4.1. *If $x \in \mathcal{P}_T$ solves (1 λ) then $V'(t, \epsilon)$ satisfies*

$$\begin{aligned} (15) \quad V'(t, \epsilon) &\leq \lambda C_s(t, t-\epsilon) \left(\int_{t-\epsilon}^t g(v, x(v))dv \right)^2 \\ &\quad + 2g(t, x) \left[\lambda C(t, t-\epsilon) \int_{t-\epsilon}^t g(v, x(v))dv - \int_{t-\epsilon}^t C(t, s)g(s, x(s))ds \right] \\ &\quad + 2g(t, x)[\lambda a(t) - x(t)]. \end{aligned}$$

Proof. Taking into account that $C_{st} \leq 0$ we have

$$\begin{aligned} V'(t, \epsilon) &\leq \lambda C_s(t, t - \epsilon) \left(\int_{t-\epsilon}^t g(v, x(v)) dv \right)^2 \\ &\quad + 2\lambda g(t, x) \int_{-\infty}^{t-\epsilon} C_s(t, s) \int_s^t g(v, x(v)) dv ds \end{aligned}$$

If we integrate the last term by parts and use (13) in the lower limiting evaluation, keeping in mind that x is bounded, we obtain

$$\begin{aligned} V'(t, \epsilon) &\leq \lambda C_s(t, t - \epsilon) \left(\int_{t-\epsilon}^t g(v, x(v)) dv \right)^2 \\ &\quad + 2\lambda g(t, x) \left[C(t, s) \int_s^t g(v, x(v)) dv \Big|_{-\infty}^{t-\epsilon} + \int_{-\infty}^{t-\epsilon} C(t, s) g(s, x(s)) ds \right] \\ &= \lambda C_s(t, t - \epsilon) \left(\int_{t-\epsilon}^t g(v, x(v)) dv \right)^2 \\ &\quad + 2\lambda g(t, x) \left[C(t, t - \epsilon) \int_{t-\epsilon}^t g(v, x(v)) dv \right] \\ &\quad + 2g(t, x) \left[\lambda \int_{-\infty}^{t-\epsilon} C(t, s) g(s, x(s)) ds + \int_{t-\epsilon}^t C(t, s) g(s, x(s)) ds \right] \\ &\quad - 2g(t, x) \int_{t-\epsilon}^t C(t, s) g(s, x(s)) ds. \end{aligned}$$

Using (1_λ) in the next-to-last term yields (15). \square

We will integrate (15) to relate $g(t, x(t))$ to $a(t)$ and then use that relation in a lower bound on the Liapunov functional to obtain the *a priori* bound. We now obtain that lower bound.

Lemma 4.2. *For any $q > 0$, if $x \in \mathcal{P}_T$ solves (1_λ) , then*

$$\begin{aligned} (x(t) - \lambda a(t))^2 &\leq 2(1 + q^{-1}) \int_{-\infty}^{t-\epsilon} C_s(t, s) ds V(t, \epsilon) \\ &\quad + 2(1 + q^{-1}) \epsilon C^2(t, t - \epsilon) \int_{t-\epsilon}^t g^2(s, x(s)) ds \\ (16) \quad &\quad + (1 + q) \left(\int_{t-\epsilon}^t |C(t, s)| ds \right)^2 \left(K \|x\| + \sup_{0 \leq u \leq T} |g(u, 0)| \right)^2. \end{aligned}$$

Proof. Let $q > 0$ be fixed and define $H = (1 + \lambda q) \left(\int_{t-\epsilon}^t C(t, s) g(s, x(s)) ds \right)^2$ so that from (1 λ) we obtain

$$\begin{aligned}
(x(t) - \lambda a(t))^2 &= \left(\lambda \int_{-\infty}^{t-\epsilon} C(t, s) g(s, x(s)) ds + \int_{t-\epsilon}^t C(t, s) g(s, x(s)) ds \right)^2 \\
&\leq \lambda(1 + q^{-1}) \left(\int_{-\infty}^{t-\epsilon} C(t, s) g(s, x(s)) ds \right)^2 + H \\
&= \lambda(1 + q^{-1}) \left(-C(t, s) \int_s^t g(u, x(u)) du \Big|_{-\infty}^{t-\epsilon} \right. \\
&\quad \left. + \int_{-\infty}^{t-\epsilon} C_s(t, s) \int_s^t g(u, x(u)) duds \right)^2 + H \\
&\text{(using (13) and } x \in \mathcal{P}_T) \\
&= \lambda(1 + q^{-1}) \left(-C(t, t - \epsilon) \int_{t-\epsilon}^t g(u, x(u)) du \right. \\
&\quad \left. + \int_{-\infty}^{t-\epsilon} C_s(t, s) \int_s^t g(u, x(u)) duds \right)^2 + H \\
&\leq 2\lambda(1 + q^{-1}) C^2(t, t - \epsilon) \left(\int_{t-\epsilon}^t g(u, x(u)) du \right)^2 \\
&\quad + 2(1 + q^{-1}) \left(\int_{-\infty}^{t-\epsilon} C_s(t, s) \int_s^t g(u, x(u)) duds \right)^2 + H \\
&\leq 2\lambda(1 + q^{-1}) C^2(t, t - \epsilon) \epsilon \int_{t-\epsilon}^t g^2(u, x(u)) du + H \\
&\quad + 2(1 + q^{-1}) \int_{-\infty}^{t-\epsilon} C_s(t, s) ds \int_{-\infty}^{t-\epsilon} C_s(t, s) \left(\int_s^t g(u, x(u)) du \right)^2 ds \\
&\leq 2\lambda(1 + q^{-1}) C^2(t, t - \epsilon) \epsilon \int_{t-\epsilon}^t g^2(u, x(u)) du \\
&\quad + 2(1 + q^{-1}) \int_{-\infty}^{t-\epsilon} C_s(t, s) ds V(t, \epsilon) \\
&\quad + (1 + q) \left(\int_{t-\epsilon}^t |C(t, s)| ds \right)^2 \left(K \|x\| + \sup_{0 \leq u \leq T} |g(u, 0)| \right)^2,
\end{aligned}$$

as required. \square

Lemma 4.3. *If*

$$(17) \quad |g(t, x)| \leq |x| \text{ for } |x| \geq L$$

where L is defined in (12), then for any $\gamma > 0$ there is an $M > 0$ such that for any solution of (1 λ) in \mathcal{P}_T we have

$$\begin{aligned}
(18) \quad V'(t, \epsilon) &\leq Ma^2(t) + [\gamma + \beta - 2]g^2(t, x(t)) + M \\
&\quad + \int_{t-\epsilon}^t [|C(t, s)| + \epsilon C_s(t, t - \epsilon) + C(t, t - \epsilon)] g^2(s, x(s)) ds.
\end{aligned}$$

Proof. By Cauchy inequality, for any $\gamma > 0$, there is an $M > 0$ such that

$$2g(t, x)a(t) \leq \gamma g^2(t, x) + Ma^2(t).$$

By (17), we may choose M so large that

$$-2g(t, x)x \leq -2g^2(t, x) + M$$

for all $t \geq 0$ and $x \in \mathfrak{R}$. Now from (15) we have

$$\begin{aligned} V'(t, \epsilon) &\leq \gamma g^2(t, x) + Ma^2(t) \\ &\quad - 2g^2(t, x) + M + C_s(t, t - \epsilon)\epsilon \int_{t-\epsilon}^t g^2(v, x(v))dv \\ &\quad + C(t, t - \epsilon) \int_{t-\epsilon}^t [g^2(t, x(t)) + g^2(v, x(v))]dv \\ &\quad + \int_{t-\epsilon}^t |C(t, s)|[g^2(t, x(t)) + g^2(s, x(s))]ds \\ &= Ma^2(t) + g^2(t, x) \left[\gamma - 2 + \epsilon C(t, t - \epsilon) + \int_{t-\epsilon}^t |C(t, s)|ds \right] + M \\ &\quad + \int_{t-\epsilon}^t [\epsilon C_s(t, t - \epsilon) + C(t, t - \epsilon) + |C(t, s)|]g^2(s, x(s))ds \\ &\text{by (9)} \\ &\leq Ma^2(t) + g^2(t, x)[\gamma + \beta - 2] + M \\ &\quad + \int_{t-\epsilon}^t [\epsilon C_s(t, t - \epsilon) + C(t, t - \epsilon) + |C(t, s)|]g^2(s, x(s))ds, \end{aligned}$$

as required. \square

Lemma 4.4. *If (17) holds, if $\epsilon \leq T$, and if γ is small enough then there is a $\mu > 0$ so that if x solves (1 $_\lambda$) and $x \in \mathcal{P}_T$ then*

$$(19) \quad \int_0^T g^2(s, x(s))ds \leq (M/\mu) \int_0^T a^2(s)ds + TM/\mu.$$

Proof. We are going to integrate (18) from 0 to T and note that $0 = V(T, \epsilon) - V(0, \epsilon)$. First, we estimate the integral of the last term in (18) as follows. We have

$$\begin{aligned} &\int_0^T \int_{t-\epsilon}^t [|C(t, s)| + \epsilon C_s(t, t - \epsilon) + C(t, t - \epsilon)]g^2(s, x(s))dsdt \\ &\leq \int_{-\epsilon}^T \int_s^{s+\epsilon} [|C(t, s)| + \epsilon C_s(t, t - \epsilon) + C(t, t - \epsilon)]dtg^2(s, x(s))ds \\ &\text{by (8)} \\ &\leq \alpha \int_{-\epsilon}^T g^2(s, x(s))ds \leq 2\alpha \int_0^T g^2(s, x(s))ds. \end{aligned}$$

With this information we now integrate (18) and obtain

$$\begin{aligned} 0 = V(T, \epsilon) - V(0, \epsilon) &\leq M \int_0^T a^2(s) ds + TM \\ &\quad + \int_0^T [\gamma - 2 + \beta + 2\alpha] g^2(s, x(s)) ds \\ &\leq M \int_0^T a^2(s) ds - \mu \int_0^T g^2(s, x(s)) ds + TM \end{aligned}$$

since $\beta + 2\alpha < 2$ and γ can be made as small as we please. \square

Lemma 4.5. *Let the conditions of Lemma 4.4 hold and suppose there is a $Q > 0$ with*

$$(20) \quad \int_{-\infty}^{t-\epsilon} C_s(t, s) (t + T - s)^2 ds \leq Q.$$

Then there is a $Q^ > 0$ with $V(t, \epsilon) \leq Q^*$.*

Proof. We have

$$\begin{aligned} V(t, \epsilon) &= \int_{-\infty}^{t-\epsilon} C_s(t, s) \left(\int_s^t g(u, x(u)) du \right)^2 ds \\ &\leq \int_{-\infty}^{t-\epsilon} C_s(t, s) (t - s) \int_s^t g^2(u, x(u)) du ds \\ &\leq \int_{-\infty}^{t-\epsilon} C_s(t, s) (t - s) \left[\int_s^{t+T} (M/\mu) a^2(u) du + (t - s + T) TM/\mu \right] ds \\ &\leq \int_{-\infty}^{t-\epsilon} C_s(t, s) (t + T - s)^2 ds [(M/\mu) \|a^2\| + TM/\mu] \end{aligned}$$

from which the result follows. \square

Lemma 4.6. *Let the conditions of Lemma 4.5 hold. Then there exists a constant $J > 0$ such that $\|x\| < J$ whenever x is T -periodic solution of (1_λ) for $0 < \lambda \leq 1$.*

Proof. By (9) and (13), we have

$$\int_{-\infty}^{t-\epsilon} C_s(t, s) ds = C(t, t - \epsilon) \leq \beta/\epsilon.$$

If $x \in \mathcal{P}_T$ solves (1_λ) , then (19) holds, and by Lemma 4.5, $V(t, \epsilon) \leq Q^*$. Now taking into account that (7) holds with $\eta < 1$, we obtain from (16) that

$$\begin{aligned} (x(t) - \lambda a(t))^2 &\leq 2(1 + q^{-1})(\beta/\epsilon)Q^* \\ &\quad + 2(1 + q^{-1})(\beta^2/\epsilon)TM(\|a^2\| + 1)/\mu \\ &\quad + (1 + q)(\eta\|x\| + \beta g^*)^2 \end{aligned}$$

where $g^* = \|g(t, 0)\|$. Since $\eta < 1$, we may choose $q > 0$ small enough so that $(1 + q)\eta^2 < 1$, and hence, there exists $J > 0$ such that $\|x\| < J$. The proof is complete. \square

5. CONTINUITY AND COMPACTNESS

We select part of (10) and define the mapping $U : \mathcal{P}_T \rightarrow \mathcal{P}_T$ by $\phi \in \mathcal{P}_T$ implies that

$$(21) \quad (U\phi)(t) = \int_{-\infty}^{t-\epsilon} C(t, s)g(s, \phi(s))ds.$$

Then U is well defined on P_T by (6). By a change of variable we have

$$(U\phi)(t) = \int_{-\infty}^t C(t, s-\epsilon)g(s-\epsilon, \phi(s-\epsilon))ds$$

with a fully convex kernel.

Lemma 5.1. *Suppose that $\int_{-\infty}^{t-\epsilon} [|C(t, s)| + |C_t(t, s)|]ds$ is bounded for all $t \in \mathfrak{R}$. Then U is continuous on P_T and for each $J > 0$, $\Gamma = \{U(\phi) : \phi \in \mathcal{P}_T, \|\phi\| \leq J\}$ is uniformly bounded and equicontinuous.*

Proof. First, there is a J^* such that $\phi \in \Gamma$ implies that $|g(t, \phi(t))| \leq J^*$ and there is a C^* with

$$(22) \quad \int_{-\infty}^{t-\epsilon} [|C(t, s)| + |C_t(t, s)|]ds \leq C^*, \quad t \in \mathfrak{R}.$$

It is clear that $U\phi \in P_T$ by (6) and the argument following (10). We now show that U is continuous on P_T . If $\tilde{\phi}, \phi \in P_T$, then

$$(23) \quad \begin{aligned} |U(\phi)(t) - U(\tilde{\phi})(t)| &= \left| \int_{-\infty}^{t-\epsilon} C(t, s)g(s, \phi(s))ds - \int_{-\infty}^{t-\epsilon} C(t, s)g(s, \tilde{\phi}(s))ds \right| \\ &= \left| \int_{-\infty}^{t-s} C(t, s) [g(s, \phi(s)) - g(s, \tilde{\phi}(s))] ds \right|. \end{aligned}$$

Since g is uniformly continuous on $[0, T] \times \{x \in R : |x| \leq \|\tilde{\phi}\| + 1\}$, for any $\epsilon > 0$, there exists $0 < \delta < 1$ such that $\|\phi - \tilde{\phi}\| < \delta$ implies $|g(s, \phi(s)) - g(s, \tilde{\phi}(s))| < \epsilon$ for all $s \in [0, T]$. It follows from (23) that $\|U(\phi) - U(\tilde{\phi})\| \leq \epsilon C^*$. Thus, F is continuous on P_T .

Next, for an arbitrary $\phi \in \Gamma$ we have

$$\frac{d}{dt}(U\phi)(t) = C(t, t-\epsilon)g(t-\epsilon, \phi(t-\epsilon)) + \int_{-\infty}^{t-\epsilon} C_t(t, s)g(s, \phi(s))ds.$$

and this derivative is bounded by

$$C(t, t-\epsilon)J^* + J^* \int_{-\infty}^{t-\epsilon} |C_t(t, s)|ds \leq J^* \sup_{0 \leq t \leq T} \|C(t, t-\epsilon)\| + J^* C^*.$$

This implies that Γ is equicontinuous. The uniform boundedness of Γ follows from the inequality

$$|U(\phi)(t)| \leq \int_{-\infty}^{t-\epsilon} |C(t, s)||g(s, \phi(s))|ds \leq J^* C^*.$$

□

6. PERIODIC SOLUTIONS

We will show the existence of T -periodic solutions of (1) by applying Theorem 2.1. By (10) and (11), we see that $x \in P_T$ is a solution of (1_λ) if and only if it is a fixed point of $B + \lambda A$.

Theorem 6.1. *If (2)-(9), (12)-(13), (17), (20), and (22) hold with $\epsilon \leq T$, then (1) has a T -periodic solution.*

Proof. Let the mappings A and B be defined in (10) and (11) with $S = P_T$. Then B is a contraction mapping with contraction constant η , and hence, $(I - B)^{-1}$ exists and is continuous on $(I - B)S = S$. By Lemma 5.1 and the Ascoli-Arzelà Theorem, we see that A is continuous and maps bounded sets into compact sets. It is also clear that $\lambda A(M) \subset (I - B)S$ for each closed convex subset $M \subset S$ and $\lambda \in [0, 1]$. Now by Lemma 4.6, the set of solutions to $x = Bx + \lambda Ax$ is bounded. Therefore, the alternative (i) of Theorem 2.1 must hold; that is, $B + A$ has a fixed point in P_T which is a T -periodic solution of (1). \square

Remark 6.2. Observe that the continuity of $C(t, s)$ with respect to s for $t - \epsilon < s < t$ is not required for fixed t . One may readily verify that the function $C(t, s)$ defined by $C(t, s) = k(t - s)^{-p}$ for $t - s \geq \epsilon$ and $C(t, s) = (t - s)^{-q}$ for $0 < t - s < \epsilon$ with $p > 2, 0 < q < 1, 0 < \epsilon \leq 1, k > 0$ satisfy all conditions of Theorem 6.1 for an appropriately chosen constant k .

REFERENCES

- [1] R. Agarwal, M. Meehan, and D. O'Regan, Fixed Point Theory and Applications, Cambridge University Press, 2001.
- [2] D. W. Boyd and J. S. W. Wong, On nonlinear contractions, Proc. Amer. Math. Soc. 20(1969), 458-464.
- [3] T. A. Burton, Integral equations, implicit functions, and fixed points, Proc. Amer. Math. Soc. 124(1996), 2383-2389.
- [4] T. A. Burton, A Liapunov functional for a singular integral equation, Nonlinear Analysis 73(2010), 3873-3882.
- [5] T. A. Burton and C. Kirk, A fixed point theorem of Krasnoselskii-Schaefer type, Math. Nachr 189(1998), 23-31.
- [6] T. A. Burton, Stability by Fixed Point Theory for Functional Differential Equations, Dover Publications, Mineola, New York, 2006.
- [7] H. Gao, Y. Li, B. Zhang, A fixed point theorem of Krasnoselskii-Schaefer type and its applications in control and periodicity of integral equations, Fixed Point Theory, To appear.
- [8] A. Granas and J. Dugundji, Fixed Point Theory, Springer-Verlag, New York, 2003.
- [9] M. A. Krasnoselskii, Some problems of nonlinear analysis, Amer. Math. Soc. Transl. 10(1958), 345-409.
- [10] Y. Liu and Z. Li, Schaefer type theorem and periodic solutions of evolution equations, J. Math. Anal. Appl. 316(2006), 237-255.
- [11] A. Meir and E. Keeler, A theorem on contractive mappings, J. Math. Anal. Appl. 28(1969), 326-329.
- [12] S. Park, Generalizations of the Krasnoselkii fixed point theorem, Nonlinear Analysis 67(2007), 3401-3410.
- [13] H. Schaefer, Über Die Methode Der a Priori Schranken, Math. Ann. 129(1955), 415-416.
- [14] D. R. Smart, Fixed Point Theorems, Cambridge University Press, Cambridge, 1980.
- [15] J. Wu, H. Xia, and B. Zhang, Topological transversality and periodic solutions of neutral functional differential equations, Proceedings of the Royal Society of Edinburgh, 129A(1999), 199-220.

- [16] B. Zhang, Liapunov functionals and periodicity in a system of nonlinear integral equations, *Electronic Journal of Qualitative Theory of Differential Equations*, Spec. Ed. I, 2009 No. 1, 1-15.